

# A likely Early Ordovician age for the regional, penetrative cleavage in the Gaissa Nappe Complex, northern Norway

BJØRN SUNDVOLL & DAVID ROBERTS

Sundvoll, B. & Roberts, D. 2003: A likely Early Ordovician age for the regional, penetrative cleavage in the Gaissa Nappe Complex, northern Norway. *Norges geologiske undersøkelse Bulletin 441*, 51–59.

Pervasively cleaved mudstones and claystones from two different parts of the Gaissa Nappe Complex in Finnmark were subjected to Rb-Sr whole-rock analytical investigation. Although neither of the two groups of samples yielded a true isochron, the data do support the concept of a probable age at around 480–500 Ma, as can be discerned from the 'isochron fits' of various combinations of the analytical data. Together with the relatively high  $^{87}\text{Sr}/^{86}\text{Sr}$  initial ratios, this result strongly suggests that a Sr-isotopic resetting event occurred at around 480–500 Ma, an event which is considered to reflect an overall, pervasive metamorphic imprint on the rock formations of the Gaissa Nappe Complex.

Earlier radiometric dating from this same allochthon has yielded ambiguous results, some pointing to Finnmarkian (Late Cambrian–Early Ordovician) deformation whereas other data have indicated a Scandian (Late Silurian–Early Devonian) age. The Rb-Sr analytical data reported here thus tend to favour a likely Finnmarkian age for the very low-grade metamorphism. As the youngest fossiliferous rocks in the Gaissa thrust sheet are of Early Tremadoc age, the cleavage-forming event is inferred to have occurred in Tremadoc or possibly Arenig time.

*Bjørn Sundvoll, Geologisk Museum, NHM, Universitet i Oslo, Postboks 1172, N-0318 Oslo, Norway.  
David Roberts, Geological Survey of Norway, N-7491 Trondheim, Norway.*

## Introduction

Rock complexes of Neoproterozoic to Early Palaeozoic age dominate the Caledonides of Finnmark, northern Norway. The nappes of the metamorphic allochthon extend over a distance of some 370 km from Kvæangen in the southwest to Varanger Peninsula in the northeast, and overlie a generally thin, parautochthonous to autochthonous sedimentary sequence that was deposited upon a peneplaned surface of Palaeoproterozoic and Archaean crystalline rocks.

The precise ages of the thick metasedimentary successions in the different parts of the Caledonide tectonostratigraphy are poorly known. Fossils are comparatively rare, occurring only in the very highest and lowest of the nappes. Likewise, our knowledge of the exact timing of Caledonian orogenic deformation is far from perfect. Although isotopic dating investigations have been made in certain areas, the data, in most cases, are not sufficiently robust to allow us to state categorically that, for example, metamorphism occurred at just one specific time. Only in the highest allochthon in this area, the Magerøy Nappe, where there are fossiliferous metasedimentary rocks of Late Ordovician to Early Silurian age, can we be sure that, in this particular case, we are dealing with *Scandian* (Silurian to Early/Mid Devonian) orogenic deformation. Elsewhere, a case has been presented for suggesting that it is possible to identify an earlier phase of Late Cambrian/Early Ordovician tectonothermal activity, known as the *Finnmarkian* (Ramsay & Sturt 1976, Sturt et al. 1978), in some of the nappes.

In this contribution, we present Rb-Sr whole-rock analytical data from the Gaissa Nappe Complex, which is a part of the Lower Allochthon of Scandinavian Caledonide tectonostratigraphy (Roberts & Gee 1985). The rocks sampled are penetratively cleaved mudstones and claystones from two widely separate areas, in southern Porsangerfjord and in the Vestertana district of innermost Tanafjord. These particular rocks were collected with the aim of trying to determine the age of this pervasive metamorphic fabric. A preliminary, unpublished report on these results has been presented earlier (Sundvoll & Roberts 2002).

## General geology

The tectonostratigraphy of the Caledonides in Finnmark has been touched upon in the introductions to several papers on diverse topics, but some of the main contributions to the overall architecture and structure of the principal nappe complexes are those of Sturt et al. (1975), Williams et al. (1976), Chapman et al. (1985), Ramsay et al. (1985), Roberts (1985), Townsend et al. (1986) and Gayer et al. (1985, 1987).

Four major nappes or nappe complexes are recognised, from bottom to top: (1) Gaissa Nappe Complex, (2) Laksefjord Nappe Complex, (3) Kalak Nappe Complex, (4) Magerøy Nappe. With the exception of the Magerøy Nappe, these various nappe complexes occur fairly widely across the mainland of Finnmark, moreso the Kalak Nappe Complex which extends southwards into the neighbouring county of Troms (Zwaan & Roberts 1978, Zwaan 1988). The




Fig. 1. Simplified tectonostratigraphy of the Caledonides of Finnmark; modified from Roberts (1985). The approximate locations of the two sampling traverses in the Gaissa Nappe Complex are shown, marked by the short, thick lines and the letters GS and SB. Details of sample locations, grid references and formation names are given in Table 2.

Magerøy Nappe occurs exclusively on the island of Magerøya in northwestern Finnmark (Fig. 1).

Briefly, the Gaissa Nappe Complex (also termed informally the Gaissa Thrust Belt, e.g. Gayer et al. 1987, Townsend et al. 1989) is dominated by low-grade, Neoproterozoic to earliest Ordovician, metasedimentary rocks, whilst the Laksefjord and Kalak Nappe Complexes are composed mainly of greenschist- to amphibolite-facies, late Mesoproterozoic to Neoproterozoic, metasedimentary successions with or without basal plinths of older Precambrian crystalline rocks (Sturt et al. 1978, Ramsay et al. 1985). Although this basement/cover contact is generally the locus of high shear strains, with mylonite development, a partly tectonised unconformity has been identified in one of the Kalak thrust sheets (Ramsay & Sturt 1977). The lithostratigraphic succession of the Kalak Nappe Complex also hosts the magmatic rocks of the extensive Seiland Igneous Province (Robins 1996). The Magerøy Nappe comprises an amphibolite-facies metasedimentary succession of Ordo-Silurian age which is cut by diverse plutonic rocks varying in composition from mafic/ultramafic to granitic (Andersen 1984, Roberts & Andersen 1985, Robins 1998).

The Magerøy Nappe and the Kalak Nappe Complex carry thick mylonites, and locally ultramylonites, along their soles. Mylonites also occur along the basal thrust to the Laksefjord Nappe Complex, but the thickness is far less. The floor thrust to the Gaissa Nappe Complex is generally characterised by a 3-5 cm-thick ultracataclasite, rarely protomylonite. Prominent stretching lineations within and above the thrust zones to all these nappes generally denote a translation sense between southeast and east-southeast. Nappe movement was polyphase in the case of the Kalak Nappe Complex, with early ductile SE translation superceded by a more brittle, ESE-directed transport (Gayer et al. 1987). A polyphase, internal deformation history is also recognised in all these nappe complexes, though in the case of the Gaissa this is much less complicated than in the other allochthons (Townsend et al. 1986).

In the Kalak Nappe Complex, there is also evidence from isotopic dating (Daly et al. 1991, Reginiussen et al. 1995, Kirkland & Daly 2003) of a pre-Caledonian, Late Riphean (>850 Ma) tectonothermal event, at least in parts of the higher thrust sheets. Whether this affects the regionally extensive, Søro Group succession in its entirety or only

partly, or is confined to basement rocks of the older Eidvågeid Sequence (Rice 1990), is a matter which has yet to be resolved. A comparable, but undated, pre-Caledonian tectonothermal event has also been reported from one of the nappes in Troms (Zwaan & van Roermund 1990).

## Gaissa Nappe Complex

Over most of its 200 km strike extent, the Gaissa Nappe Complex lies tectonically above the thin, Vendian to Lower Cambrian, condensed sequence of the Dividal Group (Føyen 1967, 1985). In the east, towards Varanger Peninsula, the precise path of the basal thrust plane is uncertain, and different solutions have been proposed (cf. Roberts 1985, Townsend 1986, Rice 1994, Sygnabere 1997). This uncertainty relates to the fact that the Gaissa and the subjacent parautochthon share a common lithostratigraphy in the southwestern part of Varanger Peninsula. In this area, and west of the Tana river, bedrock exposure is also comparatively poor.

Between innermost Laksefjord and Langfjord, in western Tanafjord, the Laksefjord Nappe Complex, or its basal, basement slice, the Kunes Nappe (Rice 2001), tectonostratigraphically overlies the Gaissa. To the southwest, the Laksefjord Nappe Complex is excised and strongly mylonitised rocks of the Kalak Nappe Complex there lie directly upon diverse formations of the Gaissa Nappe Complex (Gayer et al. 1987, Roberts 1998).

Table 1. Lithostratigraphy of the Riphean to Lower Ordovician (Tremadoc) succession occurring within the Gaissa Nappe Complex, Finnmark. Division into formations for the Digermulen, Vestertana and Tanafjorden Groups is based largely on Banks et al. (1971), Siedlecka & Siedlecki (1971) and Williams (1976). Subdivisions of the Airoaivi Group are from Townsend et al. (1989). See also the 1:250,000 map-sheets 'Vadsø' (Siedlecki 1980) and 'Honningsvåg' (Roberts 1998); and the 1:500,000 bedrock map-sheet of Finnmark county (Siedlecka & Roberts 1996).

Group	Formation	Biostratigraphic age
Digermulen	Berlogaissa	Lower Tremadoc
	Kistedalen	Middle/Upper Cambrian
	Duolbasgaissa	Lower Cambrian
Vestertana	Breidvika	Lower Cambrian
	Stappogiedde	
	Mortensnes	Vendian
	Nyborg Smalfjord	
Tanafjorden	Porsanger	Upper Riphean
	Stabbursdal	
	Hanglecærro	
	Vagge	
	Gamasfjellet	
	Dakkovarre	
	Stangenes Grønneset	
Airoaivi	Åbbardasras'sa	
	Nav'kaoaivi	
	Adnevarri	
	Bal'dneras'sa	

The lithostratigraphy of the Gaissa allochthon is shown in Table 1. Formations of the Late Riphean *Tanafjorden Group* occur throughout the nappe complex, but are more extensively exposed in the southwest. The oldest unit, the Airoaivi Group (Townsend et al. 1989), is exposed in the southwesternmost part of the nappe, west of Lakselv. The Vendian to basal Cambrian *Vestertana Group* is present from south of Laksefjord northeastwards to the southwestern parts of Varanger Peninsula, whereas the youngest unit, the Cambrian to Lower Tremadoc *Digermul Group* (Reading 1965, Banks et al. 1971, Nikolaisen & Henningsmoen 1985), occurs only on the Digermul Peninsula.

Although the various formations of these groups can be followed over distances of 200 km or more, there are facies variations and other changes which have given rise to a slightly different stratigraphic nomenclature in parts of the succession from Porsangerfjord to Varanger (Siedlecki 1980, Rice & Townsend 1996, Roberts 1998). In the Porsangerfjord district, for example, the Porsanger Formation at the top of the Tanafjorden Group overlies the Stabbursdal Formation. On Varanger, the upper member of the Grasdal Formation is equivalent to the Porsanger Formation, and the lower member of the Grasdal correlates with the Stabbursdal.

In southwestern areas, in the Porsangerfjord district, the Gaissa Nappe Complex is composed of a series of westerly dipping, imbricate thrust sheets, separated into four structural zones, including two duplexes (Townsend et al. 1986). N-S to NNE-SSW-trending folds with a steep, west-dipping, axial planar cleavage are quite common (Williams 1979). Metamorphic grade is generally in anchizone (Rice et al. 1989) but reaches up to epizone just beneath the overlying basal Kalak thrust. A somewhat simpler structural picture obtains some 200 km farther to the northeast in the inner Tanafjord-SW Varanger Peninsula region. There, imbrication is far less evident but mesoscopic folds are still common, with axes trending between N-S and NE-SW. The ESE-verging folds are open to close, asymmetric and carry a pervasive axial-planar cleavage of mainly anchizone metamorphic grade (Williams 1979, Townsend et al. 1986, Rice et al. 1989). In interbeds of mudstone type in most parts of the Gaissa Nappe Complex, the cleavage can best be described as an incipient slaty cleavage.

## Rb-Sr dating Sampling and lithologies

With the purpose of trying to determine the metamorphic age of the rocks of the Gaissa Nappe Complex, sampling was carried out in two areas: (1) along the main road between inner Vestertana and the sole thrust to the Laksefjord Nappe Complex, east of Ifjord, over a distance of roughly 20 kilometres; (2) along the road and coast north and south of Stabbursnes, western and inner part of Porsangerfjord, a distance of 12-13 km (Fig. 1). Sampling was concentrated on the most pelitic and thus better cleaved rock types in several formations. Mesoscopic and large-scale folds are ubiquitous




Fig. 2. (a) Tightly spaced cleavage in pelitic beds in multilayered sandstone and claystone/mudstone of the Nyborg Formation, at the road-cut localities GS.1-3 (see Table 2). The steeply west-dipping cleavage is axial planar to common, open to tight, c. N-S-trending folds. Photo taken by Christophe Pascal, looking south. (b) Typical character of the pervasive cleavage, axial planar to prominent folds, in interlayered maroon-violet mudstone, greenish-grey claystone and grey siltstone of the Stabburdal Formation, western shore of Porsangerfjord, near localities SB.1 and 2 (Table 2); looking approximately north-northeast. Disaggregation structures in the upper part of the photo signify that the clastic framework of the sediment has lost cohesion at some stage during its deformation history, possibly involving partial liquefaction and tectonic dewatering in the very early phase of the fold- and cleavage-forming event.

in both areas, and the cleavage is axial planar to these folds (Fig. 2).

Table 2. Sample numbers, formation names, rock characteristics and map/grid references. All map-sheets are the latest available editions, with blue coordinates. Samples GS.1-5, 1:50,000 map-sheet 2235 I 'Smalfjord', 3-NOR edition, printed 10/94; samples GS.7-15, 1:50,000 map-sheet 2235 IV 'Ifjordfjellet', 3-NOR edition, printed 7/91; samples SB.1-9, 1:50,000 map-sheet 2035 III 'Lakselv', 4-NOR edition, printed 2/94.

Sample Number	Formation	Characteristics	Map reference
GS.1	Nyborg	Green-grey slaty mudstone	2965 1510
GS.2	Nyborg	Green-grey slaty mud/claystone	2960 1510
GS.3	Nyborg	Green-grey slaty mudstone	2955 1510
GS.4	Mortensnes	Grey, cleaved tillite matrix	2970 1510
GS.5	Nyborg	Red/maroon slaty mudstone	2650 1455
GS.7	Stappogjedde	Grey-green, cleaved claystone	2135 1350
GS.10	Breidvik	Red/maroon, cleaved mudstone	1600 1665
GS.12	Breidvik	Red/maroon, cleaved mudstone	1020 1575
GS.13	Breidvik	Green, cleaved mud/claystone	1020 1575
GS.15	Breidvik	Green-grey, slate/phyllite	0730 1630
SB.1	Stabbursdal	Maroon, cleaved mudstone	2460 9465
SB.2	Stabbursdal	Grey, cleaved claystone	2460 9465
SB.3	Stabbursdal	Red.brown, cleaved mudstone	2405 9425
SB.5	Stabbursdal	Grey/maroon, cleaved mudstone	2210 9305
SB.7	Stabbursdal	Maroon, cleaved mudstone	2030 8765
SB.8	Stabbursdal	Dark grey/black, cleaved mudstone	2085 8765
SB.9	Dakkovarre	Dark grey, cleaved claystone	2075 8295

A list of the samples used in this study, their formation affiliations, general features and grid references is given in Table 2. Detailed geology at 1:50,000 scale is available only for map-sheet 'Lakselv' (Roberts & Davidsen 1992). For a more regional compilation of the geology the reader is referred to the 1:250,000 map-sheet 'Honningsvåg' (Roberts 1998).

### Analytical techniques

Rb and Sr concentrations were determined by isotope dilution technique using a combined spike of enriched <sup>87</sup>Rb and <sup>84</sup>Sr, as described by Boelrijk (1968). The uncertainty of the method is estimated to 0.5% in the Rb/Sr ratios. All isotopic measurements were carried out on a VG354 TIMS instrument at the Mineralogical-Geological Museum, University of Oslo, following methods similar to those described by Jacobsen & Heier (1978). During the period of analysis the NBS987 Sr-isotopic standard yielded an average of 0.71024 ± 3. Isochron calculations were carried out according to the method described by York (1969). The calculations were based on the value of 1.42 x 10<sup>-11</sup> a<sup>-1</sup> for the decay of <sup>87</sup>Rb (Steiger & Jäger 1977). All results are given as 2σ errors.




Fig. 3. The analysed samples plotted on a  $^{87}\text{Sr}/^{86}\text{Sr}$  versus  $^{87}\text{Rb}/^{86}\text{Sr}$  diagram. (a) The sample series 'GS', from the northeastern part of the Gaissa Nappe Complex (cf. Fig. 1). (b) The sample series 'SB' from the western part of the Gaissa Nappe Complex. The analytical data and isochron calculations are shown in Tables 3 and 4, respectively.

## Results

The results of the elemental and isotopic determinations and the calculated Rb/Sr and corresponding  $^{87}\text{Rb}/^{86}\text{Sr}$  ratios are presented in Table 3. The results of the isochron calculations are listed in Table 4. The data are also plotted in Figs. 3 a & b.

As can be seen from Table 4, cases 1 and 3, neither of the two groups of samples yielded any true isochron, i.e., with a MSWD of <2.5. There are no clear-cut indications or reasons for leaving any one or more samples out of the calculations. The semi-linear relationship of the data as seen in Figs. 3 a & b, probably reflects random secondary effects on the Rb/Sr system in the mudstone and claystone rocks. However, the close and true isochron ages indicated by cases 2, 4 and 5 (Table 4), and the relatively high  $^{87}\text{Sr}/^{86}\text{Sr}$  initial ratios (0.732–0.735), suggest an event of Sr-isotopic resetting at around 480–500 Ma, which is thought to reflect an overall, pervasive metamorphic imprint on the rock formations from which the samples are taken.

## Discussion

### Gaissa Nappe Complex

Although not definitive, the results of this Rb-Sr analytical investigation strongly suggest that resetting of the Sr iso-

Table 3. Rb and Sr analytical data. (Rb and Sr concentration errors approx. 1.5%).

Sample	Rb(ppm)	Sr(ppm)	Rb/Sr	$^{87}\text{Rb}/^{86}\text{Sr}$	$^{87}\text{Sr}/^{86}\text{Sr}$	SE
GS1	243.49	27.66	8.084	25.980	.91125	±.00003
GS2	170.60	31.00	5.505	16.141	.84400	±.00003
GS3	204.73	29.70	6.893	20.246	.86354	±.00005
GS4	142.79	112.83	1.266	3.680	.75885	±.00003
GS5	218.95	65.44	3.346	9.768	.80013	±.00003
GS7	165.95	42.11	3.941	11.493	.79015	±.00003
GS10	124.33	38.41	3.237	9.440	.79055	±.00003
GS12	213.07	48.35	4.407	12.893	.82193	±.00003
GS13	150.41	25.85	5.820	17.094	.86324	±.00003
GS15	129.99	27.17	4.784	14.007	.83016	±.00003
SB1	258.40	20.36	12.690	37.722	.98833	±.00003
SB2	233.00	30.56	7.625	22.471	.89746	±.00003
SB3	218.06	53.10	4.107	12.014	.82162	±.00003
SB5	186.61	65.10	2.866	8.364	.79487	±.00003
SB7	184.61	83.87	2.201	6.065	.78028	±.00003
SB8	224.07	81.21	2.759	8.065	.81390	±.00003
SB9	148.65	100.17	1.484	4.318	.76521	±.00003

Table 4. Isochron calculations.

Case	Samples	Age error	$(^{87}\text{Sr}/^{86}\text{Sr})_i$	error	MSWD
1	all GS samples	480 ± 60	.7319	±.007	125
2	all except GS3, GS7, GS13	475 ± 32	.7334	±.004	25
3	all SB samples	548 ± 128	.7339	±.014	226
4	all except SB8	495 ± 22	.7351	±.002	9
5	all except SB1, SB8	512 ± 10	.7336	±.001	.1

$(^{87}\text{Sr}/^{86}\text{Sr})_i$  = calculated initial ratio of  $^{87}\text{Sr}/^{86}\text{Sr}$

tope system occurred at around 480-500 Ma. Moreover, the very high initial  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio further suggests that this resetting can, in all probability, be ascribed to a pervasive metamorphic imprint on the rock formations of the Tanafjorden and Vestertana Groups. This calls for an assessment of earlier isotopic studies from this particular nappe complex.

The earliest radiometric work on the Gaissa is that of Pringle (1973; and in Sturt et al. 1975) who reported a Rb-Sr whole-rock isochron age of  $504 \pm 7$  Ma (recalculated to a 1.42 decay constant), interpreted as the age of cleavage formation in pelites of the Stappogiedde Formation (Vestertana Group). This 'isochron' was subsequently rejected by Dallmeyer & Reuter (1989) and Dallmeyer et al. (1989) on the grounds that complete isotopic equilibration was probably not achieved, and thus detrital Rb-Sr systems were not fully rejuvenated.

This study was followed by that of Dallmeyer (1988b), who reported on the  $^{40}\text{Ar}/^{39}\text{Ar}$  geochronology of whole-rock, cleaved pelite samples from four undetermined locations within the Gaissa Nappe Complex. He noted that spectra of samples taken from the hinge zones of cleavage-related folds gave «well-defined plateaux...which correspond to ages of 474 and 482 Ma.» These ages were interpreted by Dallmeyer (1988b) to approximately date cleavage formation at these locations, a metamorphic event which this same author considered was probably related to late-stage Finnmarkian tectonothermal activity.

Subsequently, Dallmeyer et al. (1989) carried out a joint  $^{40}\text{Ar}/^{39}\text{Ar}$  and K-Ar study on five samples of penetratively cleaved pelites from Gaissa formations, and then came to a different conclusion, namely that the earliest metamorphism had a maximum age of c. 440 Ma. Although whole-rock,  $^{40}\text{Ar}/^{39}\text{Ar}$ , incremental-release, total-gas ages for the five samples ranged between c. 474 and 550 Ma, more reliance was placed on one sample taken from the southwestern part of the Gaissa Nappe Complex, in Porsangerfjord, where the <0.5 mm size fraction of illite gave a K-Ar apparent age of  $440.2 \pm 9.4$  Ma. In this same paper (Dallmeyer et al. 1989), another apparent age of  $463.6 \pm 9.9$  Ma, and also in the <0.5 mm size-fraction range, from the northeastern part of the Gaissa Nappe Complex, was not discussed.

Preference for the one K/Ar apparent age at the expense of the other K-Ar and  $^{40}\text{Ar}/^{39}\text{Ar}$  data was influenced by the work of Reuter (1987) in the Rheinisches Schiefergebirge in Germany, who had demonstrated that only the very smallest size fraction yielded ages that were compatible with the geological constraints in that area. Coarser size fractions were found to retain a detrital memory, and thus gave imprecise, 'mixed' ages. Dallmeyer et al. (1989) therefore preferred to reject the  $^{40}\text{Ar}/^{39}\text{Ar}$  analyses; and, at the same time, Dallmeyer's (1988b) own earlier work and conclusions from rocks of the Gaissa Nappe Complex were also rejected. They also argued that an Early Silurian metamorphic age was more consistent with «that of the Scandian orogenesis recorded within late Proterozoic through Early Silurian

autochthonous/-parautochthonous metasedimentary sequences exposed in the central and southern Scandinavian Caledonides.» However, this contention, that low-grade metamorphism and deformation of Scandian age within the Lower Allochthon in southern areas must also be Scandian 1000 km farther north, is clearly open to debate.

In a palaeomagnetic study of low-grade sedimentary rocks from the parautochthon and Gaissa Nappe Complex of southwestern Varanger Peninsula, Bylund (1994) observed that two of three samples from sites within the Gaissa along the shores of Tanafjord yielded data which indicated that a remagnetization had occurred in these rocks in Early Ordovician time. The implications of these observations were not discussed further.

Field evidence from basal parts of the Gaissa Nappe Complex shows that thrusting post-dated the peak of metamorphism and cleavage generation. One sample of ultracataclasite from the sole thrust (from Handelsbukta, c. 12 km ENE of Lakselv) has been dated by the K-Ar method, and has given an age of  $391 \pm 9$  Ma (J.G. Mitchell, unpubl. data, pers. comm. 1989; reported in Roberts & Sundvoll 1990, with permission). This Mid Devonian age is comparable to dates generated from top-ESE shear bands affecting mylonites at the base of the Kalak Nappe Complex (see below).

### Laksefjord and Kalak Nappe Complexes

In order to fully appreciate the data presented in this contribution, it is relevant to consider some of the published isotopic data from the suprajacent Laksefjord and Kalak Nappe Complexes.

The earliest attempt at dating the penetrative slaty cleavage in pelites of the Friarfjord Formation, *Laksefjord Nappe Complex*, is that of Pringle in Sturt et al. (1978). Samples taken from the slate quarry at Friarfjord generated a Rb-Sr whole-rock isochron age of  $492 \pm 45$  Ma (recalculated using a 1.42 decay constant), which was interpreted to date the peak of metamorphism. The rocks of the Laksefjord Nappe Complex were thus considered by Sturt et al. (1978) to have been involved in Finnmarkian orogenic deformation. Subsequently, Dallmeyer et al. (1989) sampled from the same slate quarry and obtained a  $^{40}\text{Ar}/^{39}\text{Ar}$  total-gas age of  $485 \pm 9$  Ma.

Several isotopic dating studies carried out within the supracrustal and magmatic rocks of the *Kalak Nappe Complex* have provided evidence for either Scandian or Finnmarkian metamorphic events, or both. Early K/Ar and Rb/Sr studies in West Finnmark (Sturt et al. 1967, Pringle & Sturt 1969) laid a basis for recognising an early-Caledonian, Finnmarkian event in rocks of the Kalak Nappe Complex (Sturt et al. 1978). Although these early data, and their interpretation, were shrouded in controversy (Krill & Zwaan 1987, Binns 1989, Townsend & Gayer 1989), subsequent and more precise  $^{40}\text{Ar}/^{39}\text{Ar}$  and Sm-Nd dating from different areas in northern Norway and in the correlative Seve Nappes of northern Sweden (Dallmeyer 1988a, Dallmeyer et al. 1988,

1991, Mørk et al. 1988, Essex et al. 1997) confirmed this important, latest Cambrian to earliest Ordovician, orogenic phase. Zwaan & van Roermund 1990) also recognised this early, amphibolite-facies and initial thrusting event in the Corrovarre Nappe of Troms and suggested possible correlations with the eclogite-bearing Seve Nappes of northern Sweden.

In an investigation involving Rb-Sr whole-rock and thin-slab dating of mylonites from the Kalak Thrust Zone near Børselv, eastern Porsangerfjord, one set of 11 samples yielded an isochron age of  $479 \pm 15$  Ma (MSWD = 2.26) (Roberts & Sundvoll 1990). In another sample series taken from close to the base of the thrust zone, two separate subsets of thin slabs of top-ESE, shear-banded mylonites gave isochron ages of  $385 \pm 29$  (MSWD = 2.51) and  $380 \pm 22$  (MSWD = 3.04) Ma, respectively. This thin-slab dating indicates resetting of the Rb-Sr isotope systems on a local scale in Mid Devonian time, broadly coeval with the latest phase of ESE-directed nappe translation. Although the  $479 \pm 15$  Ma isochron date could be taken to indicate ductile mylonite generation at around the transition from Tremadoc to Arenig time, Roberts & Sundvoll (1990) were careful to point out possible complications related to the Rb-Sr isotope system with these particular samples.

In northwestern Varanger Peninsula, a  $^{40}\text{Ar}/^{39}\text{Ar}$  study on the penetrative cleavage in lower to mid greenschist-facies phyllites of the Berlevåg Formation in the Tanahorn Nappe, considered to represent a part of the Kalak Nappe Complex (Levell & Roberts 1977), has yielded an age of c. 460 Ma (Frank & Rice 1999, Rice & Frank, in press). This metamorphism and deformation is regarded by these authors as representing a late stage of the Finnmarkian tectonothermal event. Samples taken from beneath the Tanahorn Nappe in formations of the Barents Sea Region (Fig. 1) gave a pre-470 Ma age for this same cleavage, whereas a later, spaced, crenulation cleavage yielded an age of c. 443 Ma (Rice & Frank, in press). There are thus clear indications that the main syn-metamorphic, SE-directed fold deformation in the Tanahorn Nappe and its footwall is pre-Scandian and of Ordovician age.

In summary, there is evidence, both in the literature and from ongoing research, that parts of the metamorphic allochthon of Finnmark were involved in early-Caledonian (Late Cambrian to Early Ordovician) orogenic deformation, particularly at the Middle and lowermost Upper Allochthon levels represented by the Laksefjord and Kalak Nappe Complexes. In the Gaissa Nappe Complex, the evidence has been ambiguous. While early radiometric work provided indications of a Finnmarkian event, Dallmeyer and coworkers rejected these data, thereby denying the existence of an Early Ordovician tectonothermal event in the metasedimentary rocks of the Gaissa Nappe Complex.

The Rb-Sr analytical data presented here therefore bring us back to square one. While we prefer to keep an open mind in this particular instance, in view of the interpretative

problems associated with the Rb-Sr whole-rock dating method, it is pertinent to note that the strong indications of a metamorphic event at around 480-500 Ma fit well with the biostratigraphic evidence from the youngest rocks in the nappe, the Digermul Group, which extend up into the Early Tremadoc. With the Cambrian-Ordovician boundary now defined at c. 490 Ma and the base Arenig at c. 480 Ma (Davidek et al. 1998, Landing et al. 2000), deformation in the Gaissa Nappe Complex could conceivably have started shortly after deposition of the Berlogaissa Formation (Reading 1965). Although paleocurrent data are lacking from most of the Digermul Group, Reading (1965) has noted the sudden incursion of scour-and-fill cross-bedding close to the exposed top of the Berlogaissa. Speculatively, this may be indicating the onset of a fluvial regime as an emergent topography developed in the west and northwest ahead of a rising Finnmarkian mountain chain. A detailed sedimentological study of the three formations of the Digermul Group would be a prerequisite for taking these ideas further.

## Conclusions

Samples of pervasively cleaved mudstones and claystones from two different parts of the Gaissa Nappe Complex in Finnmark were subjected to Rb-Sr analytical investigation with the aim of trying to determine the metamorphic age of the rocks in this thrust sheet. Although neither of the two groups of samples yielded a true isochron, the data as a whole do support the concept of a probable isochron age at around 480-500 Ma, as can be discerned from the 'isochron fits' of various combinations of the analytical data. Together with the relatively high  $^{87}\text{Sr}/^{86}\text{Sr}$  initial ratios, this result strongly suggests that a Sr-isotopic resetting event occurred at around 480-500 Ma. This event is thought to reflect an overall, pervasive metamorphic imprint on the rock formations of the Gaissa Nappe Complex.

Previous radiometric dating evidence from the Gaissa Nappe Complex has given ambiguous results, some pointing to Finnmarkian deformation whereas other data have indicated a Scandian age. The Rb-Sr analytical data reported here do tend to favour a likely Finnmarkian age for the very low-grade metamorphism; and this fits with diverse isotopic data obtained from the overlying nappe complexes, as well as with the biostratigraphic evidence where the youngest rocks in the Gaissa Nappe Complex are of Early Tremadoc age.

## Acknowledgements

We are most grateful to Professors Donald Ramsay and Torgeir Andersen for their helpful and constructive reviews of the manuscript. Thoughtful and important suggestions by Professor Arne Råheim and Dr. Bouke Zwaan also led to improvements in the final typescript. Dr. Christophe Pascal kindly provided a replacement field photograph for Figure 2a.

## References

- Andersen, T.B. 1984: The stratigraphy of the Magerøy Supergroup, Finnmark, North Norway. *Norges geologiske undersøkelse Bulletin* 395, 25-37.
- Banks, N.L., Edwards, M.B., Geddes, W.P., Hobday, D.K. & Reading, H.G. 1971: Late Precambrian and Cambro-Ordovician sedimentation in East Finnmark. *Norges geologiske undersøkelse* 269, 197-236.
- Binns, R.E. 1989: Regional correlations in NE Troms-W. Finnmark: the demise of the 'Finnmarkian' orogeny? In Gayer, R.A. (ed.) *The Caledonide geology of Scandinavia*, Graham & Trotman, London, 27-45.
- Boelrijk, N.A.I.M. 1968: A general formula for 'double' isotope dilution analysis. *Chemical Geology* 3, 323-325.
- Bylund, G. 1994: Palaeomagnetism of Vendian-Early Cambrian sedimentary rocks from E.Finnmark, Norway. *Tectonophysics* 231, 45-57.
- Dallmeyer, R.D. 1988a: Polyorogenic  $^{40}\text{Ar}/^{39}\text{Ar}$  mineral age record within the Kalak Nappe Complex, northern Scandinavian Caledonides. *Journal of the Geological Society, London* 145, 705-716.
- Dallmeyer, R.D. 1988b: Polyphase tectonothermal evolution of the Scandinavian Caledonides. In Harris, A.L. & Fettes, D.J. (eds.) *The Caledonian-Appalachian orogen*. Geological Society Special Publication 38, 365-379.
- Dallmeyer, R.D. & Reuter, A. 1989:  $^{40}\text{Ar}/^{39}\text{Ar}$  whole-rock dating and the age of cleavage in the Finnmark autochthon, northernmost Scandinavian Caledonides. *Lithos* 22, 213-227.
- Dallmeyer, R.D., Mitchell, J.G., Pharaoh, T.C., Reuter, A. & Andersen, A. 1988: K-Ar and  $^{40}\text{Ar}/^{39}\text{Ar}$  whole-rock ages of slate/phyllite from autochthonous-parautochthonous sequences in the Komagfjord and Alta-Kvænangen tectonic windows, northern Scandinavian Caledonides: evaluating the extent and timing of Caledonian tectonothermal activity. *Bulletin of the Geological Society of America* 100, 1493-1501.
- Dallmeyer, R.D., Reuter, A., Clauer, N. & Liewig, N. 1989: Chronology of Caledonian tectonothermal activity within the Gaissa and Laksefjord Nappe Complexes (Lower Allochthon), Finnmark, Norway: evidence from K-Ar and  $^{40}\text{Ar}/^{39}\text{Ar}$  ages. In Gayer, R.A. (ed.) *The Caledonide geology of Scandinavia*. Graham & Trotman, London, 9-26.
- Dallmeyer, R.D., Andréasson, P.G. & Svenningsen, O. 1991: Initial tectonothermal evolution within the Scandinavian Caledonide accretionary prism: constraints from  $^{40}\text{Ar}/^{39}\text{Ar}$  mineral ages within the Seve Nappe Complex, Sarek Mountains, Sweden. *Journal of Metamorphic Geology* 9, 203-218.
- Daly, J.S., Aitchison, S.J., Cliff, R.A., Gayer, R.A. & Rice, A.H.N. 1991: Geochronological evidence from discordant plutons for a late Proterozoic orogen in the Caledonides of Finnmark, northern Norway. *Journal of the Geological Society, London* 148, 29-40.
- Davidek, K., Landing, E., Bowring, S.A., Westrop, S.R., Rushton, A.W.A., Fortey, R.A. & Adrain, T.M. 1998: New uppermost Cambrian U-Pb date from Avalonian Wales and age of the Cambrian-Ordovician boundary. *Geological Magazine* 135, 305-309.
- Essex, R.M., Gromet, L.P., Andréasson, P.G. & Albrecht, L. 1997: Early Ordovician U-Pb metamorphic ages of the eclogite-bearing Seve Nappes, northern Scandinavian Caledonides. *Journal of Metamorphic Geology* 15, 665-676.
- Frank, W. & Rice, A.H.N. 1999:  $^{40}\text{Ar}/^{39}\text{Ar}$  cleavage age data from the North Varanger Region, Finnmark, N. Norway, and its bearing on Caledonian-Baikalian interaction. (abstract). 10<sup>th</sup> EUG Congress, Strasbourg, Abstract Volume, 93-94.
- Føyn, S. 1967: Dividal-gruppen (Hylolithus-sonen) i Finnmark og dens forhold til de eokambrikk-kambriske formasjoner. *Norges geologiske undersøkelse* 249, 1-82.
- Føyn, S. 1985: The Late Precambrian in northern Scandinavia. In Gee, D.G. & Sturt, B.A. (eds.) *The Caledonide orogen - Scandinavia and related areas*. John Wiley & Sons, Chichester, 233-245.
- Gayer, R.A., Hayes, S.J. & Rice, A.H.N. 1985: The structural development of the Kalak Nappe Complex of eastern and central Porsangerhalvøya, Finnmark, Norway. *Norges geologiske undersøkelse Bulletin* 400, 67-87.
- Gayer, R.A., Rice, A.H.N., Roberts, D., Townsend, C. & Welbon, A. 1987: Restoration of the Caledonian Baltoscandian margin from balanced cross-sections: the problem of excess continental crust. *Transactions of the Royal Society of Edinburgh: Earth Sciences* 78, 197-217.
- Jacobsen, S.B. & Heier, K.S. 1978: Rb/Sr isotope systematics in metamorphic rocks, Kongsberg sector, South Norway. *Lithos* 11, 257-276.
- Kirkland, C.L. & Daly, J.S. 2003: Confirmation of the Porsanger orogeny in the Caledonides of Arctic Norway. (Abstract) *British Geological Survey Highlands Workshop*, Edinburgh, May 2003, Abstract volume.
- Krill, A.G. & Zwaan, K.B. 1987: Reinterpretation of Finnmarkian deformation on western Sørøy, northern Norway. *Norsk Geologisk Tidsskrift* 67, 15-24.
- Landing, E., Bowring, A.S., Davidek, K.L., Rushton, A.W.A., Fortey, R.A. & Wimbledon, W.A.P. 2000: Cambrian-Ordovician boundary age and duration of the lowest Ordovician Tremadoc Series based on U-Pb zircon dates from Avalonian Wales. *Geological Magazine* 137, 485-494.
- Levell, B.K. & Roberts, D. 1977: A re-investigation of the geology of north-west Varanger Peninsula, east Finnmark, north Norway. *Norges geologiske undersøkelse* 334, 83-90.
- Mørk, M.B.E., Kullerud, K. & Stabell, A. 1988: Sm-Nd dating of Seve eclogites, Norbotten, Sweden - evidence for early Caledonian (505 Ma) subduction. *Contributions to Mineralogy and Petrology* 99, 344-351.
- Nikolaisen, F. & Henningsmoen, G. 1985: Upper Cambrian and lower Tremadoc olenid trilobites from the Digermul Peninsula, northern Norway. *Norges geologiske undersøkelse Bulletin* 400, 1-49.
- Pringle, I.R. 1973: Rb-Sr age determinations on shales associated with the Varanger Ice Age. *Geological Magazine* 109, 465-472.
- Pringle, I.R. & Sturt, B.A. 1969: The age of the peak of the Caledonian orogeny in west Finnmark, northern Norway. *Norsk Geologisk Tidsskrift* 49, 435-436.
- Ramsay, D.M. & Sturt, B.A. 1976: The syn-metamorphic emplacement of the Magerøy Nappe. *Norsk Geologisk Tidsskrift* 56, 291-307.
- Ramsay, D.M. & Sturt, B.A. 1977: A sub-Caledonian unconformity within the Finnmarkian nappe sequence and its regional significance. *Norges geologiske undersøkelse* 334, 107-116.
- Ramsay, D.M., Sturt, B.A., Zwaan, K.B. & Roberts, D. 1985: The Caledonides of northern Norway. In Gee, D.G. & Sturt, B.A. (eds.) *The Caledonide orogen - Scandinavia and related areas*. John Wiley & Sons, Chichester, 163-184.
- Reading, H.G. 1965: Eocambrian and Lower Palaeozoic geology of the Digermul Peninsula, Tanafjord, Finnmark. *Norges geologiske undersøkelse* 234, 167-191.
- Reginiussen, H., Krogh Ravna, E.G. & Berglund, K. 1995: Mafic dykes from Øksfjord, Seiland Igneous Province, northern Norway: geochemistry and palaeotectonic significance. *Geological Magazine* 132, 667-681.
- Reuter, A. 1987: Implications of K-Ar ages of whole-rock and grain-size fractions of metapelites and intercalated metatuffs within an anchizonal terrane. *Contributions to Mineralogy and Petrology* 97, 105-115.
- Rice, A.H.N. 1994: Stratigraphic overlap of the Late Proterozoic Vadsø and Barents Sea Groups and correlation across the Trollfjorden-Komagelva Fault Zone, Finnmark, North Norway. *Norsk Geologisk Tidsskrift* 74, 48-57.
- Rice, A.H.N. 2001: Field evidence for thrusting of the basement rocks coring tectonic windows in the Scandinavian Caledonides; an insight from the Kunes Nappe, Finnmark, Norway. *Norsk Geologisk Tidsskrift* 81, 321-328.
- Rice, A.H.N. & Townsend, C. 1996: Correlation of the Late Precambrian Ekkerøya Formation (Vadsø Group; E. Finnmark) and the Brennelvfjord Interbedded Member (Porsangerfjord Group; W. Finnmark), N. Norwegian Caledonides. *Norsk Geologisk Tidsskrift* 76, 55-61.



- Rice, A.H.N. & Frank, W. 2003: The early Caledonian (Finnmarkian) event reassessed in Finnmark:  $^{40}\text{Ar}/^{39}\text{Ar}$  cleavage age data from NW Varangerhalvøya, N.Norway. *Tectonophysics*, in press.
- Rice, A.H.N., Bevins, R.E., Robinson, D. & Roberts, D. 1989: Evolution of low-grade metamorphic zones in the Caledonides of Finnmark, North Norway. In Gayer, R.A. (ed.) *The Caledonide geology of Scandinavia*. Graham & Trotman, London, 172-192.
- Roberts, D. 1985: The Caledonian fold belt in Finnmark: a synopsis. *Norges geologiske undersøkelse Bulletin 403*, 161-177.
- Roberts, D. 1998: Berggrunnskart Honningsvåg – Geologisk kart over Norge, M 1:250 000. *Norges geologiske undersøkelse*.
- Roberts, D. & Andersen, T.B. 1985: Nordkapp. Beskrivelse til det berggrunnsgeologiske kartbladet, M 1:250 000. *Norges geologiske undersøkelse Skrifter 61*, 1-49.
- Roberts, D. & Gee, D.G. 1985: An introduction to the structure of the Scandinavian Caledonides. In Gee, D.G. & Sturt, B.A. (eds.) *The Caledonide orogen – Scandinavia and related areas*. John Wiley & Sons, Chichester, 55-68.
- Roberts, D. & Sundvoll, B. 1990: Rb-Sr whole-rock and thin-slab dating of mylonites from the Kalak Thrust Zone, near Børselv, Finnmark. *Norsk Geologisk Tidsskrift 70*, 259-266.
- Roberts, D. & Davidsen, B. 1992: Lakselv: berggrunnsgeologisk kart 2035 3, M 1:500 000, foreløpig utgave. *Norges geologiske undersøkelse*.
- Robins, B. 1996: The Seiland Igneous Province, North Norway: field trip guidebook, IGCP Project 336. *Norges geologiske undersøkelse Report 96.127*, 34 pp.
- Robins, B. 1998: The mode of emplacement of the Honningsvåg Intrusive Suite, Magerøya, northern Norway. *Geological Magazine 135*, 231-244.
- Siedlecka, A. & Siedlecki, S. 1971: Late Precambrian sedimentary rocks of the Tanafjord-Varangerfjord Region of Varanger Peninsula, northern Norway. *Norges geologiske undersøkelse 269*, 246-294.
- Siedlecka, A. & Roberts, D. 1996: Finnmark fylke: Berggrunnsgeologi M 1:500 000. *Norges geologiske undersøkelse*.
- Siedlecki, S. 1980: Geologisk kart over Norge, berggrunnskart Vadsø – M 1:250 000. *Norges geologiske undersøkelse*.
- Steiger, K.H. & Jäger, E. 1977: Subcommission on geochronology: convention on use of decay constants in geo- and cosmochronology. *Earth and Planetary Science Letters 36*, 359-362.
- Sturt, B.A., Miller, J.A. & Fitch, F.J. 1967: The age of alkaline rocks from West Finnmark, northern Norway, and their bearing on the dating of the Caledonian Orogeny. *Norsk Geologisk Tidsskrift 47*, 255-273.
- Sturt, B.A., Pringle, I.R. & Roberts, D. 1975: Caledonian nappe sequence of Finnmark, northern Norway, and the timing of orogenic deformation and metamorphism. *Bulletin of the Geological Society of America 86*, 710-718.
- Sturt, B.A., Pringle, I.R. & Ramsay, D.M. 1978: The Finnmarkian Phase of the Caledonian Orogeny. *Journal of the Geological Society, London 135*, 597-610.
- Sundvoll, B. & Roberts, D. 2003: Rb-Sr dating of cleaved mudstones from eastern and western parts of the Gaissa Nappe Complex, Finnmark. *Norges geologiske undersøkelse Report 2002.110*, 20 pp.
- Sygnabere, T.O. 1997: *Strukturgeologisk kartlegging av den kaledonske fronten i de østlige delene av Tanafjord-Varangerfjordregionen, Varangerhalvøya (Finnmark), og dens betydning for utviklingen av Trollfjord-Komagelvforkastningssonen*. Cand. Scient. thesis, University of Bergen, 127 pp.
- Townsend, C. 1986: *Thrust tectonics within the Caledonides of northern Norway*. Unpubl. Ph.D. thesis, University of Wales, Cardiff, 137 pp.
- Townsend, C., Roberts, D., Rice, A.H.N. & Gayer, R.A. 1986: The Gaissa Nappe, Finnmark, North Norway: an example of a deeply eroded imbricate zone within the Scandinavian Caledonides. *Journal of Structural Geology 8*, 431-440.
- Townsend, C., Rice, A.H.N. & Mackay, A. 1989: The structure and stratigraphy of the southwestern portion of the Gaissa Thrust Belt and the adjacent Kalak Nappe Complex, Finnmark, N.Norway. In Gayer, R.A. (ed.) *The Caledonide geology of Scandinavia*. Graham & Trotman, London, 111-126.
- Williams, D.M. 1976: A revised stratigraphy of the Gaissa Nappe, Finnmark. *Norges geologiske undersøkelse 324*, 63-78.
- Williams, D.M. 1979: Structural development of the Gaissa Nappe in the Finnmark Caledonides, North Norway. *Norges geologiske undersøkelse 348*, 93-104.
- Williams, G.D., Rhodes, S., Powell, D.B., Pässe, C.R., Noake, J.S. & Gayer, R.A. 1976: A revised tectono-stratigraphy for the Kalak Nappe in Central Finnmark. *Norges geologiske undersøkelse 324*, 47-61.
- York, D. 1969: Least squares fitting of a straight line with correlated errors. *Earth and Planetary Science Letters 5*, 320-324.
- Zwaan, K.B. 1988: Nordreisa, berggrunnsgeologisk kart – M 1:250 000. *Norges geologiske undersøkelse*.
- Zwaan, K.B. & Roberts, D. 1978: Tectonostratigraphic succession and development of the Finnmarkian nappe sequence, North Norway. *Norges geologiske undersøkelse 343*, 53-71.
- Zwaan, K.B. & van Roermund, H.L.M. 1990: A rift-related mafic dyke swarm in the Corrovarre Nappe of the Caledonian Middle Allochthon, Troms, North Norway, and its tectonometamorphic evolution. *Norges geologiske undersøkelse Bulletin 419*, 25-44.